

RECOMMENDATIONS



order@tigersul.com

www.tigersul.com

TIGER 90CR® sulphur

— Granular Sulphur —

Product Description

TIGER 90CR[®] **sulphur** is a high analysis granular degradable sulphur product that can be used both as a source of plant nutrient sulphur and / or as a soil amendment for correction of problem alkali soils. Yearly applications of TIGER 90CR[®] will improve soil sulphur levels as well as loss associated with nutrients that are soluble.

TIGER 90CR® sulphur is a unique, high analysis sulphur fertilizer that delivers agronomically and economically, and provides excellent handling characteristics.

Guaranteed Analysis - 90% Sulphur		
Physical Properties		
Color	Pale Copper Pastilles	
Fertilizer Prill Size	Regular Grade - 260 SGN	Mini Prill Grade - 190 SGN
Bulk Density (packed)	Regular Grade - 76 lbs/ft³	Mini Prill Grade - 88 lbs/ft²

General Application and Use Recommendations

Use **TIGER 90CR**[®] **sulphur** (0-0-0-90) on *Every acre of every crop in every year* to build soil sulphur levels. (including forages and hay land).

For 1 pound of sulphur apply 1.11 lbs of TIGER 90CR® sulphur.

TIGER 90CR[®] **sulphur**, when used as a part of a balanced fertility program, can provide a season-long source of sulphur. **TIGER 90CR**[®] **sulphur** can be applied alone or blended with granular fertilizers. Amount required should be based on agronomic recommendations from soil tests.

Recommendations should also be based on crop removal rates of sulphur.

TIGER 90CR® Sulphur Management Strategies

To start or convert to a **TIGER 90CR**[®] **sulphur maintenance program**, especially on low sulphur soils or for high sulphur-demand crops, **follow one of three programs:**

- 1. Supplement **TIGER 90CR[®] Organic Sulphur** with additional soluble sulphate (20-0-0-24, 19-2-0-22, etc) fertilizer in the first year, or;
- 2. Start a **TIGER 90CR[®] Organic Sulphur** maintenance program on fields where a low sulphur demand crop is being grown and then apply yearly to all crops, including forages, OR;

TIGER 90CR® Sulphur Recommendations

3. Surface apply **TIGER 90CR**[®] **sulphur** in the fall to allow early granule dispersion and measure soil sulphate the following spring to determine the need for supplemental sulphate fertilizer the first year of a maintenance program.

If **TIGER 90CR**[®] **sulphur** is not applied regularly as part of a fertilizer program, or if soil sulphur levels are deficient, then soluble sulphate should be applied to high sulphur-using crops such as oil seeds and legumes. *Under no circumstances should* **TIGER 90CR**[®] **sulphur** *be* applied alone to sulphur deficient soil unless the application rate applied is adequate enough to overcome the sulphur deficiency. In northern soils, Tiger-Sul Products recommends an application of a combination of sulphate and sulphur to assist with the building of sulphur reserves in these soils. Consult an agronomic specialist in these rare cases.

For more information on TIGER 90CR® sulphur view the "TIGER 90CR Sulphur Advantage" in our Sulphur Technical information.

Soil Amending with TIGER 90CR® Sulphur — Applications of 300 to 1000 lbs are common for soil amendments. For more information view the "TIGER 90CR® sulphur Soil Amending" technical sheet.

Handling and Storage

TIGER 90CR® Sulphur will blend and store well with other granular fertilizer products. Accepted blending and handling procedures should be followed when using TIGER 90CR® Sulphur. Good housekeeping is necessary when handling sulphur products. Tiger-Sul Products uses a special anti-dust agent to minimize air borne dust. Always use stringent dust control procedures to prevent concentrated flammable dusts from reaching a spark or flame source. Sulphur materials should not be blended and stored with strong oxidizing agents such as ammonium nitrate. Avoid extensive use of augers due to fracturing of the product.

First Aid

Skin Contact: Wash with mild soap and water.

Eyes: Irrigate thoroughly with copious quantities of plain water. Inadequate irrigation may cause irritation. Do not use Boric Acid.

Ingestion: Solid Sulphur is virtually non-toxic. It can be taken internally at fairly large doses without injury.

Inhalation: Sulphur dust may irritate the mucous membranes of the respiratory passages.

*Disclaimer: While every reasonable care has been taken to ensure that uniformity on our web site is correct at the time of posting, we do not warrant accuracy and recommend you obtain local and independent verification.



